

## JALUPRO AFTERCARE

The following aftercare advice is essential to help reduce the risk of treatment complications, to improve your comfort/healing and to help achieve the best possible results from treatment.

Please follow the aftercare advice for Jalupro injections.

- Immediately after Jalupro injections the treatment area may be red, tender or swollen. There may be red raised bumps in the injection sites, these effects will usually be resolved by 2 days.
- For significant swelling contact your practitioner for advice. Swelling may be reduced by regular cold compress, anti-inflammatory medicines such as Ibuprofen (after 24 hours, if safe for you to take), antihistamines and sleeping with your head propped up on 3 pillows to help drainage of fluid overnight.
- Bruising may occur following treatment, bruising will resolve naturally but may take up to 2 weeks to settle.
- Regular application of Arnica cream may be helpful for bruising.
- Avoid any makeup for 24 hours after treatment as this will reduce the risk of infection.
- Drink plenty of water after treatment to keep well hydrated.
- Avoid alcohol and caffeine for 24 hours after treatment.
- Avoid medications such as aspirin or Ibuprofen for 24 hours as they are blood thinning agents and may worsen bruising (if these medications have been advised by your GP you must check with your GP before stopping them)
- Avoid supplements such as multivitamins, fish oils, glucosamine and Vitamin B or Vitamin E, which are blood thinning agents and may exacerbate bruising for 24 hours (if these supplements have been advised by your GP please check with your GP before stopping them)
- Avoid any excess sun exposure, extremes of hot/cold, saunas, swimming or strenuous exercise until the redness and swelling has settled.
- You must seek medical attention and contact your practitioner if you experience any signs or symptoms of infection after treatment. Infection can present as hot, red shiny skin, there may be pus formation and you may have a fever or feel generally unwell.
- You must seek emergency medical attention if you experience any severe allergy symptoms after treatment. Symptoms may include rash, facial swelling and breathing difficulties.
- Contact your practitioner as soon as possible if you notice any other unwanted side effects.
- Your practitioner will advise when further treatment appointments are required. If you are advised to attend a follow up appointment, please do make every effort to attend them. You should do this even if you believe that  
the recovery process is going well and you cannot see that there are any visible complications.

It is essential that you have read all of the information available. Once you have read and understood all of the below, please sign the declaration at the bottom of this form. Please do let your practitioner know if you have any questions or if you do not understand any of the aftercare instructions provided below.